

## Low Latitude Ionospheric Scintillation and Zonal Plasma Irregularity Drifts Climatology around the Equatorial Anomaly Crest Over Kenya

O.J. Olwendo<sup>1</sup>; P. Baki\*<sup>2</sup>; P.J. Cilliers<sup>3</sup>; P. Doherty<sup>4</sup> & S. Radicella<sup>5</sup>

<sup>1</sup>Pwani University College; \*<sup>2</sup>Department of Physics, Technical University of Kenya; <sup>3</sup>South African National Space Agency, Space Science Directorate; <sup>4</sup>Institute for Scientific Research, Boston College; <sup>5</sup>The Abdus Salam International Centre for Theoretical Physics, Italy

### Abstract

In this study we have used a VHF and GPS-SCINDA receiver located at Nairobi (36.8°E, 1.3°S, dip -24.1°) in Kenya to investigate the climatology of ionospheric L-band scintillation occurrences for the period 2009 to 2012; and seasonal variation of the zonal plasma drift irregularities derived from a VHF receiver for the period 2011. The annual and diurnal variations of L-band scintillation indicate occurrence at post sunset hours and peaks in the equinoctial months. However VHF scintillation occurs at all seasons around the year and is characterized by longer duration of activity and a slow fading that continues till early morning hours unlike in the L-band where they cease after midnight hours. A directional analysis has shown that the spatial distribution of scintillation events is mainly on the Southern and Western part of the sky over Nairobi station closer to the edges of the crest of the Equatorial Ionization Anomaly. The distribution of zonal drift velocities of the VHF related scintillation structures indicates that they move at velocities in the range of 20–160 m/s and their dimension in the East–West direction is in the range of 100–00 km. The December solstice is associated with the largest plasma bubbles in the range of 600–900 km. The most significant observation from this study is the occurrence of post-midnight scintillation without pre-midnight scintillations during magnetically quiet periods. The mechanism leading to the formation of the plasma density irregularity causing scintillation is believed to be via the Rayleigh Tailor Instability; it is however not clear whether we can also attribute the post-midnight plasma bubbles during magnetic quiet times to the same mechanism. From our observations in this study, we suggest that a more likely cause of the east ward zonal electric fields at post-midnight hours is the coupling of the ionosphere with the lower atmosphere during nighttime. This however needs a further investigation based on relevant data.

### Keywords

- Ionospheric scintillation;
- Zonal drifts velocity;
- GPS-SCINDA;
- VHF receiver

*Journal of Atmospheric and Solar-Terrestrial Physics, Vol.138–139 pp, 9-22. (2016).*

See more at: <http://www.sciencedirect.com/science/article/pii/S136468261530105X>